

University of Groningen

N4Py iron complexes as model for iron bleomycin

Roelfes, J.G.; Hage, R.; Que, L.; Feringa, B.L.

Published in:
Journal of Inorganic Biochemistry

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1999

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Roelfes, J. G., Hage, R., Que, L., & Feringa, B. L. (1999). N4Py iron complexes as model for iron bleomycin. *Journal of Inorganic Biochemistry*, 74(8), 279 - 279.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Spectroscopic studies of FixL, an O₂ sensing heme protein

Kenton R. Rodgers, Gudrun S. Lukat-Rodgers, Lei Tang & Lisa Volk
 Department of Chemistry, North Dakota State University, Fargo, ND 58105
 rogers@plains.nodak.edu

FixL is the sensor component of the bacterial two-component signal transduction system responsible for regulating N₂ fixation in several species of bacteria. FixL contains a heme domain capable of reversible O₂ binding and a kinase domain whose activity is inhibited in response to heme oxygenation. This inhibition results in failure to phosphorylate the response regulator, FixJ, leaving it transcriptionally inactive toward *nifA* and *fixK*. We are pursuing a molecular-level understanding of the mechanism by which FixL's kinase activity is coupled to binding or release of O₂ by the heme. Data from cw and time-resolved spectroscopic methods will be presented and interpreted in terms of a model for transmission of the signal between the heme and kinase domains of FixL.



N4Py iron complexes as model for iron bleomycin

Gerard Roelfes, Ronald Hage, Lawrence Que, Jr. and Ben L. Feringa

University of Groningen, Nijenborgh 4, 9747 AG Groningen, the Netherlands; Unilever Research Laboratories, Olivier van Noortlaan 120, 3133 AT Vlaardingen, the Netherlands; University of Minnesota, 55455 MN Minneapolis, USA.
 Roelfes@chem.rug.nl

Iron-peroxo species are invoked in the mechanisms of a number of iron requiring biological oxidation catalysts. For example a low spin Fe(III)-OOH species has been characterized for "activated Bleomycin", the active form of the antitumor drug bleomycin (BLM). Activated BLM is formed by the reaction of Fe(II)-BLM with O₂ in the presence of a 1e⁻ reductant. The decomposition of this intermediate is thought to be responsible for the drug's ability to oxidatively cleave DNA. The accepted mechanistic proposal involves hydrogen abstraction by activated BLM at the C4' position of the deoxyribose unit of a nucleotide to form a C-4' carbon radical whose fate is determined by two subsequent pathways, one that requires additional O₂ and another that does not. Besides DNA cleavage activated BLM is also capable of oxidizing a wide variety of organic substrates.

In our efforts to gain a better understanding of the chemistry of iron Bleomycin we prepared a series of model complexes from the pentadentate ligand N4Py. Complexation of this ligand with iron(II) or iron(III) salts leads to the formation of a series of stable and well characterized complexes including [(N4Py)Fe(CH₃CN)](ClO₄)₂ (1), [(N4Py)FeOMe](ClO₄)₂ (2) and [{N4Py)Fe}₂O](ClO₄)₂ (3). Reaction of H₂O₂ lead to the formation of a transient purple species which has been spectroscopically (UV/Vis, EPR and ES-MS) characterized to be [(N4Py)Fe(III)OOH]²⁺ (4). Resonance Raman spectroscopy showed the presence of weakened O-O bond. The purple intermediate 4 is formed from 1 in a two step process: first oxidation to give two iron(III) species, 2 and 3, followed by an acid base reaction to give 4. The reactivity of 1 in catalytic oxidations has been studied. A wide variety of organic substrates, including alkanes, alkenes, alcohols and aromatic substrates could be oxidized by 1, using H₂O₂ as the oxidant. Monitoring the catalytic reactions with UV/Vis showed that 4 is involved in the catalytic reaction as the precursor for the active species. A possible mechanism for oxidations catalyzed by 1 will be discussed.

